



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

parabola with reference to its axis are  $\frac{a \cos^3 \theta}{\cos^2 \frac{1}{2} \theta}$  and  $\frac{2a \sin \theta \cos^2 \theta}{\cos^2 \frac{1}{2} \theta}$ , and its parameter  $\frac{a \sin^2 \theta \cos \theta}{\cos^2 \frac{1}{2} \theta}$ .

Denoting by  $x$  and  $y$  the current co-ordinates of this curve with reference to the axis of the given parabola, and the focus as origin, we find from the above

$$x = \frac{a \cos \theta + a \sin^2 \theta \cos \theta - a \cos^3 \theta}{\cos^2 \frac{1}{2} \theta} = \frac{a \sin 2\theta \sin \theta}{\cos^2 \frac{1}{2} \theta},$$

$$y = a \frac{\sin \theta - \sin 2\theta \cos \theta}{\cos^2 \frac{1}{2} \theta} = -\frac{a \sin \theta \cos 2\theta}{\cos^2 \frac{1}{2} \theta}.$$

By eliminating  $\theta$  we get  $(x^2 + y^2 - 4a^2)^2 (x^2 + y^2) = y^2 (x^2 + y^2 + 4a^2)^2$ . Introducing polar co-ordinates  $\rho$  and  $\phi$ , we get the simpler equation

$$\rho = \pm 2a \cot(45^\circ - \frac{1}{2}\phi) \text{ and } \rho = \pm 2a \tan(45^\circ - \frac{1}{2}\phi).$$

Also solved by G. B. M. Zerr.

#### AVERAGE AND PROBABILITY.

160. Proposed by J. F. LAWRENCE, A. M., Stillwater, Oklahoma.

Two points are taken at random in a triangle, the line joining them dividing the triangle into two portions. Find the mean value of that portion containing the center of gravity.

Solution by HENRY HEATON, Belfield, N. D.

The triangle may be considered equilateral (see Williamson's *Integral Calculus*, p. 355). Put  $\triangle$  = area of equilateral triangle whose side =  $a$ . Let  $P$  be one of the points,  $PD = y$  and  $BD = x$ . Let  $EPF$  be the line through the two points, and put  $\angle BFP = \theta$  and  $\angle BEP = \phi$ . Then  $\phi = \frac{2}{3}\pi - \theta$ ,  $PF = y \operatorname{cosec} \theta$  and  $DF = y \cot \theta$ . The area of the elemental triangle  $PFF' = \frac{1}{2} y^2 \operatorname{cosec}^2 \theta d\theta$ . We will suppose the second point to be confined to this elemental triangle. Put  $BF = z$ . Then  $z = x + y \cot \theta$ ,  $BE = z \sin \theta \operatorname{cosec} \phi$ , and area of triangle  $EBF = \frac{\triangle z^2}{a^2} \sin \theta \operatorname{cosec} \phi$ .

If  $\theta < \frac{1}{6}\pi$  and  $F$  is confined to the line  $BC$  the area of the portion of the triangle containing the center of gravity is  $\frac{\triangle}{a^2} (a^2 - z^2 \sin \theta \operatorname{cosec} \theta)$ . The limits of  $z$  for this are  $\frac{2}{3}y \sqrt{3} \sin \phi \operatorname{cosec} \theta$  and  $a$ . Those of  $y$  are 0 and  $\frac{1}{2}a \sqrt{3} \sin \theta \operatorname{cosec} \phi$ . If  $\theta > \frac{1}{6}\pi$  and  $< \frac{1}{3}\pi$ , the line  $EF$  passes through  $O$  when  $z = \frac{a}{3} (1 + \sin \phi \operatorname{cosec} \theta)$ .

If  $z$  is less than this,  $y$  is less than  $\frac{a}{6}\sqrt{3}(1+\sin\theta \operatorname{cosec}\phi)$ .

If  $z > \frac{a}{3}(1+\sin\phi \operatorname{cosec}\theta)$  and  $y < \frac{a}{6}\sqrt{3}(1+\sin\theta \operatorname{cosec}\phi)$  the limits of  $z$  are  $\frac{a}{3}(1+\sin\phi \operatorname{cosec}\theta)$  and  $a$ .

$$\begin{aligned} \Delta = \Delta \left[ \int_0^{\frac{1}{2}\pi} \frac{1}{3^2} \sin\theta \operatorname{cosec}^3\phi - \frac{1}{4^8} \sin^2\theta \operatorname{cosec}^4\phi + \int_{\frac{1}{2}\pi}^{\frac{3}{2}\pi} \frac{1}{4^8} \sin^2\theta \operatorname{cosec}^4\phi - \frac{1}{2^7} \frac{1}{3^7} \right. \\ \times (\sin^2\theta \operatorname{cosec}^4\phi + \sin^2\phi \operatorname{cosec}^2\theta) + \frac{1}{2^5} \frac{1}{3^6} (\sin\theta \operatorname{cosec}^3\phi + \sin\phi \operatorname{cosec}^3\theta + \frac{41}{2^4} \frac{1}{3^7} \\ \left. \times \operatorname{cosec}\theta \operatorname{cosec}\phi \right] d\theta \div \frac{1}{3^2} \int_0^{\frac{1}{2}\pi} \sin\theta \operatorname{cosec}^3\phi d\theta = \frac{\Delta}{3^6} (470 + \frac{4}{3} \log 2) = .6997 \Delta. \end{aligned}$$

This is problem 76, p. 513, Williamson's *Integral Calculus*.

170. Proposed by LON C. WALKER, A. M., Santa Barbara, California.

Find the area of a triangle formed by drawing a line at random through each of three points taken at random within a given triangle.

Solution by G. B. M. ZERR, Ph. D., Parsons, W. Va.


Let  $ABC$  be the given triangle;  $P, Q, R$  the random points;  $AI=h$ ,  $BI=d$ ,  $CI=e$ ,  $d+e=a$ ,  $AM=u$ ,  $MR=v$ ,  $AS=w$ ,  $SP=z$ ,  $AN=m$ ,  $NQ=n$ .  $y-v=r(x-u)$ , the line through  $R$ ... (1),  $y-z=s(x-w)$ , the line through  $P$ ... (2),  $y-n=t(x-m)$ , the line through  $Q$ ... (3), where  $r=\tan\theta$ ,  $s=\tan\phi$ ,  $t=\tan\psi$ .

The intersection of (1) and (2) is given by

$$x_1 = \frac{ru - ws + z - v}{r - s}, \quad y_1 = \frac{rsu - rsw + rz - sv}{r - s}.$$

The intersection of (1) and (3) is given by

$$x_2 = \frac{ru - mt + n - v}{r - t}, \quad y_2 = \frac{rtu - mrt + rn - tv}{r - t}.$$



The intersection of (2) and (3) is given by

$$x_3 = \frac{sw - mt + n - z}{s - t}, \quad y_3 = \frac{stw - mst + sn - tz}{s - t}.$$

$$\begin{aligned} \text{Area of triangle} &= \frac{1}{2}(x_2y_1 - x_1y_2 + x_3y_2 - x_2y_3 + x_1y_3 - x_3y_1) \\ &= \frac{1}{2}[(ur-v)(s-t) + (ws-z)(t-r) + (mt-n)(r-s)]^2 / (r-s)(r-t)(s-t) \\ &= A/B. \end{aligned}$$

The limits of  $u$  are 0 and  $h$ ; of  $w$ , 0 and  $u$ ; of  $m$ ,  $w$  and  $u$ ; of  $v$ ,  $-du/h=v_1$  and  $eu/h=v_2$ ; of  $z$ ,  $-dw/h=w_1$  and  $ew/h=w_2$ ; of  $n$ ,  $-dm/h=m_1$  and  $em/h=m_2$ . The number of ways the three points can be taken on the surface of the triangle is  $\frac{1}{8}(\frac{1}{2}ah)^3 = \frac{1}{8}a^3h^3$ .